

S. J. Surma, A. Wroński, S. Zachorowski

ON JAŚKOWSKI – TYPE SEMANTICS FOR THE INTUITIONISTIC PROPOSITIONAL LOGIC

This is a lecture read at the XX Conference for the History of Logic, Cracow, April 1974.

One of remarkable results of Jaśkowski is the construction of a sequence of finite matrices adequate for the intuitionistic propositional logic. This result was published in 1936 in the paper [2] where only a very condensed sketch of proof is to be found. It was 17 years before a more detailed proof was published in [4] by Rose who worked out some modification of the strategy suggested by Jaśkowski for eluding the lemma which he was unable to prove. A detailed proof following closely the Jaśkowski's strategy is presented in [6]. The sequence of finite matrices adequate for the intuitionistic propositional logic (*INT*) was obtained by Jaśkowski as a result of alternate application the operation of direct power and s.c. Γ -operation to the two-element Boolean algebra. This Γ -operation of Jaśkowski can be considered as a special case of the sum operation for pseudo-Boolean algebras introduced later by Troelstra [7]. Suppose we are given the pseudo-Boolean algebras \mathcal{A} and \mathcal{B} with the sum $\mathcal{A} \oplus \mathcal{B}$ is the pseudo-Boolean algebra with the universe $A \cup B$ and the lattice ordering $\leq_{\mathcal{A} \oplus \mathcal{B}} = \leq_{\mathcal{A}} \cup \leq_{\mathcal{B}} \cup (A \times B)$. The result of Jaśkowski's Γ -operation performed on the pseudo-Boolean algebra \mathcal{A} is isomorphic to $\mathcal{A} \oplus \mathcal{H}$ where \mathcal{H} is the two-element Boolean algebra. Thus, the diagram of the lattice ordering of $\Gamma(\mathcal{A})$ can be obtained from that of \mathcal{A} by adding the new greatest element. The sequence $\{\mathcal{F}_n : n = 1, 2, \dots\}$ constructed in [2] is given by the conditions: $\mathcal{F}_1 = \mathcal{H}$, $\mathcal{F}_{n+1} = \Gamma(\mathcal{F}_n)$. Denoting the content of \mathcal{F}_n by $E(\mathcal{F}_n)$ one can express the main result of Jaśkowski as follows:

THEOREM. $\bigcap (E(\mathcal{F}_n) : n = 1, 2, \dots) = INT$.

It should be mentioned that the theorem above immediately yields finite approximability of INT .

Let N be the set of positive integers. By Jaśkowski-type sequence determined by a mapping $f : N \rightarrow N$ and a pseudo-Boolean algebra \mathcal{A} we mean the sequence $\mathcal{A}^f = \{\mathcal{A}_n^f : n \in N\}$ such that $\mathcal{A}_1^f = \mathcal{A}$, $\mathcal{A}_{n+1}^f = \Gamma((\mathcal{A}_n^f)^{f(n)})$. Let us note the following generalization of Jaśkowski's theorem:

PROPOSITION 1. *Let \mathcal{A}^f be a Jaśkowski-type sequence. If the algebra \mathcal{A} is finite then the following conditions are equivalent:*

- (i) $\bigcap (E(\mathcal{A}_n^f) : n \in N) = INT$;
- (ii) for every $n \in N$ there exists $m \in N$ such that $n < f(m)$.

Note that Jaśkowski's theorem follows by Proposition 1 and the observation that the sequence $\{\mathcal{F}_n : n \in N\}$ can be described as \mathcal{H}^Δ where Δ is the identity mapping of N . Proposition 1 shows that it is impossible to improve the result of Jaśkowski in the direction suggested by McKay [3] who stated erroneously that the sequence \mathcal{H}^f such that $f(n) = 2$ for every $n \in N$ is adequate for INT .


Suppose we are given a sequence of mappings $F = \{f^n : n \in N\}$ such that $f^n : N \rightarrow N$ for every $n \in N$. By diagonal sequence determined by F and a pseudo-Boolean algebra \mathcal{A} we mean the sequence $\mathcal{A}^F = \{\mathcal{A}_n^F : n \in N\}$ such that for every $n \in N$, \mathcal{A}_n^F is the n -th algebra of the Jaśkowski-type sequence \mathcal{A}^{f^n} (i.e. $\mathcal{A}_n^F = \mathcal{A}_{f^n(n)}^F$). Let us note the following fact about diagonal sequences (it can be generalized in various ways):

PROPOSITION 2. *Let \mathcal{A}^F be a diagonal sequence. If $n < m$ implies that $f^n(i) < f^m(i)$ for every $n, m, i \in N$ then $\bigcap (E(\mathcal{A}_n^F) : n \in N) = INT$.*

By the propositions stated here to construct a sequence of pseudo-Boolean algebras adequate for INT is very easy. In particular, one can obtain a simple sequence ϑ^Δ where ϑ is the degenerate pseudo-Boolean algebra and a diagonal sequence \mathcal{H}^C where $C = \{c^n : n \in N\}$ consists of the constant mappings : $N \rightarrow N$ $c^n(i) = n$ for every $n, i \in N$.

It is known (see [1]) that every finite pseudo-Boolean algebra is determined (up to isomorphism) by the ordered set of join-irreducible elements. Moreover, the ordered set of join-irreducible elements considered as a Kripke-frame (see [5]) has the content equal to that of the algebra. Thus, having a sequence of finite pseudo-Boolean algebras adequate for INT one can construct the corresponding sequence of Kripke-frames which

also is adequate. The sequences of frames corresponding to \mathcal{H}^Δ , ϑ^Δ and \mathcal{H}^C are described in Smoryński [5] where a very elegant Kripke-style proof of Jaśkowski's theorem is to be found. The frames corresponding to the successive algebras of the sequences \mathcal{H}^Δ , ϑ^Δ and \mathcal{H}^C can be visualized by means of the following diagrams (the empty frame corresponds to the degenerate algebra ϑ):



References

- [1] F. Birkhoff, **Lattice Theory**, Third Edition, Providence, Rhode Island, 1967.
- [2] S. Jaśkowski, *Recherches sur le système de la logique intuitioniste*, **Actualities sci. industrielles** 393 (1936), pp. 58–61.
- [3] C. G. McKay, *A note on Jaśkowski sequence*, **Zeitschrift Math. Log. und Grund. Math** 13 (1967), pp. 95–96.
- [4] F. G. Rose, *Propositional calculus and realizability*, **Trans. Amer. Math. Soc.** 75 (1953), pp. 1–19.
- [5] C. Smoryński, **Investigations of intuitionistic formal systems by means of Kripke models**, Doctoral dissertation, Stanford Univ.,

1972.

[6] S. J. Surma, *Jaśkowski's matrix criterion for the intuitionistic propositional calculus*, **Studies in the Hist. of. Math. Logic**, Cracow, 1973, pp. 87–121.

[7] A. S. Troelstra, *On intermediate propositional logics*, **Indag. Math.** 27 (1965), pp. 141–152.

Department of Logic
Jagiellonian University
Cracow