

Hiroakira Ono

## ON SOME INTUITIONISTIC MODAL LOGICS

This is an abstract of my paper *On some intuitionistic modal logics* submitted to **Publ. RIMS, Kyoto Univ.**.

Some modal logics based on logics weaker than the classical logic have been studied by Fitch [4], Prior [7], Bull [1], [2], [3], Prawitz [6] etc. Here we treat modal logics based on the intuitionistic propositional logic, which call intuitionistic modal logics (abbreviated as IML's).

Let  $H$  be the intuitionistic propositional logic formulated in the Hilbert-style. The rules of inference of  $H$  are modus ponens and the rule of substitution. The IML  $L_0$  is obtained from  $H$  by adding the following three axioms,

$$\begin{aligned}\Box p &\supset p, \\ \Box p &\supset \Box\Box p, \\ \Box(p \supset q) &\supset (\Box p \supset \Box q),\end{aligned}$$

and the rule of necessitation, i.e., from  $A$  infer  $\Box A$ . It is clear that  $L_0$  with the law of excluded middle becomes  $S4$ . Now, consider the following axioms.

$$\begin{aligned}A_1 &: \neg\Box p \supset \Box\neg\Box p, \\ A_2 &: (\Box p \supset \Box q) \supset \Box(\Box p \supset \Box q), \\ A_3 &: \Box(\Box p \vee q) \supset (\Box p \vee \Box q), \\ A_4 &: \Box p \vee \Box\neg\Box p.\end{aligned}$$

The logic  $L_0$  with the axiom  $A_i$  is denoted by  $L_i$  for  $i = 1, 2, 3, 4$ . The logic  $L_3$  with  $A_1$  (or  $A_2$ ) is denoted by  $L_{31}$  (or  $L_{32}$ ). It is easy to see that  $S4$  with any  $A_i$  is equal to  $S5$ .

We identity a logic  $L$  with the set of formulas provable in  $L$ .

THEOREM 1.

- (i) For  $J = 1, 2, 3, 31, 32$ ,  $L_0 \subsetneq L_J \subsetneq L_4$ .
- (ii)  $L_1 \subsetneq L_2 \subsetneq L_{32}$  and  $L_3 \subsetneq L_{31} \subsetneq L_{32}$ .

For IML's, we introduce a kind of Kripke models, which we call  $I$  models. A triple  $(M, \leq, R)$  is an  $I$  frame, if

- (i)  $M$  is a nonempty set with a partial order  $\leq$ ,
- (ii)  $R$  is a reflexive and transitive relation on  $M$  such that  $x \leq y$  implies  $xRy$  each  $x, y \in M$ .

For any formula  $A$  and an element  $a \in M$ , a valuation  $W(A, a) \in \{t, f\}$  is defined in the same way as a valuation on a Kripke model for the intuitionistic propositional logic. For instance,

$$W(A \supset B, a) = T \text{ if and only if for any } b \text{ such that } a \leq b, W(A, b) = f \text{ or } W(B, b) = t.$$

Moreover, we claim that

$$W(\Box A, a) = t \text{ if and only if for any } b \text{ such that } aRb, W(A, b) = t.$$

A quadruple  $(M, \leq, R, W)$  is an  $I$  model if  $(M, \leq, R)$  is an  $I$  frame and  $W$  is a valuation on it. A formula  $A$  is valid in an  $I$  frame  $(M, \leq, R)$  if  $W(A, a) = t$  for any valuation  $W$  on  $(M, \leq, R)$  and any element  $a \in M$ .

For any binary relation  $R$ , we write  $x \sim_R y$  if  $xRy$  and  $yRx$  hold. In what follows we omit the subscript letter  $R$ . Now define  $I$  frames of type  $J$  for  $J = 0, 1, 2, 3, 31, 32, 4$  as follows.

- (0) Any  $I$  frame is of type 0.
- (1) An  $I$  frame  $(M, \leq, R)$  is of type 1 when for each  $x, y \in M$ , if  $xRy$  then there is an element  $y'$  in  $M$  such that  $x \leq y'$  and  $yRy'$ .
- (2) An  $I$  frame  $(M, \leq, R)$  is type 2 when for each  $x, y \in M$ , if  $xRy$  then there is an element  $y'$  in  $M$  such that  $x \leq y'$  and  $y \sim y'$ .
- (3) An  $I$  frame  $(M, \leq, R)$  is of type 3 when for each  $x, y \in M$ , if  $xRy$  then there is an element  $x'$  in  $M$  such that  $x \sim x'$  and  $x' \leq y$ .
- (3i) An  $I$  frame is of type  $3i$  if it is both of type 3 and of type  $i$ , for  $i = 1, 2$ .
- (4) An  $I$  frame  $(M, \leq, R)$  is of type 4 if  $R$  is symmetric.

THEOREM 2. *A formula is provable in  $L_J$  if and only if it is valid in any  $I$  frame of type  $J$ , for  $J = 0, 1, 2, 3, 31, 32, 4$ .*

An IML  $L_J$  has the finite model property if for any formula  $A$  not provable in  $L_J$  there is a finite  $I$  frame of type  $J$  in which  $A$  is not valid.

THEOREM 3. *For  $J = 0, 2, 3, 32, 4$ ,  $L_J$  has the finite model property.*

In [5], another kind of Kripke models is introduced and discussed.

## References

- [1] R. A. Bull, *A modal extension of intuitionistic logic*, **Notre Dame Journal of Formal Logic** 6 (1965), pp. 142–146.
- [2] R. A. Bull, *Some modal calculi based on IC*, **Formal systems and recursive functions**, ed. by J. N. Crossley and M. A. E. Dummett, Amsterdam, 1965, pp. 3–7.
- [3] R. A. Bull, *MIPC as the formalisation of an intuitionist concept of modality*, **Journal of Symbolic Logic** 31 (1966), pp. 609–616.
- [4] F. B. Fitch, *Intuitionistic modal logic with quantifiers*, **Portugaliae Mathematica** 7 (1948), pp. 113–118.
- [5] H. Ono, *On some intuitionistic modal logics*, to appear in **Publications of the Research Institute for Mathematical Sciences**, Kyoto University.
- [6] D. Prawitz, *Natural deduction, A proof theoretical study*, **Stockholm studies in Philosophy** 3, Stockholm, 1965.
- [7] A. N. Prior, **Time and modality**, Oxford, 1957.

*College of Integrated Arts and Sciences  
Hiroshima University, Hiroshima*